

Gamma Function

by Thomas Yamada

February 4, 2026

What is the Gamma Function?

The Gamma Function is defined as:

$$\Gamma(z) = \int_{u=0}^{\infty} u^{z-1} e^{-u} dt \quad (1)$$

Graph of $\Gamma(z)$:




Figure 1: Graph of the Gamma Function $\Gamma(z)$.

You can see, by the graph, that the function is neither odd or even, because its asymmetric everywhere. There is no linearity either.

There is two relations though. It's core identity is the recurrence relation, the Gamma's equivalent of a "linearity rule":

$$\Gamma(z + 1) = z\Gamma(z) \quad (2)$$

Then, you have Euler's Reflection Formula, a sort of complementary behavior:

$$\Gamma(z)\Gamma(1 - z) = \frac{\pi}{\sin(\pi z)} \quad (3)$$

So, what is this Gamma Function really? Remember the factorial? Well,

$$\Gamma(z+1) = z! \quad (4)$$

It's partially that simple, but also not exactly. If you remember using the factorial, you've only ever used integers, because that's easy to compute with the basic function. i.e. $6! = 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = 720$

But, how would you do that with $\frac{1}{2}!$? What about $-\frac{1}{2}!$? That's where the Gamma Function steps in. It allows you to compute the "factorial" of non integers. It does have some restrictions though:

$$\text{Dom}(\Gamma) = \mathbb{R} \setminus \{-n : n \in \mathbb{Z}_{\geq 0}\} \quad (5)$$

Let's try $\Gamma(1)$:

$$\begin{aligned} \Gamma(1) &= \int_{u=0}^{\infty} u^0 e^{-u} du = \int_{u=0}^{\infty} e^{-u} du \\ \Gamma(1) &= \left[-e^{-u} \right]_{u=0}^{\infty} = 0 - (-1) \\ \Gamma(1) &= 1 = 0! \end{aligned} \quad (6)$$

For $\Gamma(2)$, you'll need to use a simple IBP:

$$\begin{aligned} \Gamma(2) &= \int_{u=0}^{\infty} u^1 e^{-u} du = \int_{u=0}^{\infty} t e^{-u} du \\ \Gamma(2) &= -ue^{-u} + \int_{u=0}^{\infty} e^{-u} du = 0 + 1 \\ \Gamma(2) &= 1 = 1! \end{aligned} \quad (7)$$

You can probably see where this is going. $\Gamma(3) = 2 = 2!$

So, what about the recurrence relation? How can we use that? Well, keep in mind that computing $\Gamma(0)$ does not exist, as you can see from the graph above. So, if we were to apply $\Gamma(z+1) = z\Gamma(z)$ to get, for example, a negative integer, you'd need to re-use the relation again and again until you reached a computable number, but you'd always hit 0 before anything actually computable, which is why every negative integer less than 0 does not exist.

Then, what about negative fractionals? Let's use some negative fractional that is perfectly divisible by $\frac{1}{2}$, such as $-\frac{3}{2}$. You'll need to use the recurrence relation in order to get to that. The first directly computable value for that is $\Gamma(\frac{1}{2})$, so we need to solve that first. Luckily, it gives a nice number to work with.

Let's look into how we'd find $\Gamma\left(\frac{1}{2}\right)$:

$$\Gamma\left(\frac{1}{2}\right) = \int_{u=0}^{\infty} e^{-u} u^{-\frac{1}{2}} du$$

Let's rewrite it so it's easier to read.

$$\Gamma\left(\frac{1}{2}\right) = \int_{u=0}^{\infty} e^{-u} \frac{1}{\sqrt{u}} du$$

Now, let's apply u -sub,

$$\text{Let: } w = \sqrt{u}, \quad dw = \frac{1}{2\sqrt{u}} du, \quad u = w^2$$

Next, substitute back in and factor out the 2.

$$\Gamma\left(\frac{1}{2}\right) = 2 \int_{w=0}^{\infty} e^{-w^2} dw$$

Looking at the graph for this new function, we can see that the integral can consume the 2 and expand the integral bounds from $0 \rightarrow \infty$ to $-\infty \rightarrow \infty$.




Figure 2: Graph of the Bell Curve $f(x) = e^{-x^2}$.

So, let's change the bounds. Let's also set the integral to some arbitrary variable, J .

$$J = \int_{w=-\infty}^{\infty} e^{-w^2} dw$$

Since e^{-w^2} has no elementary anti derivative, we can use a Multivariable Calculus trick to solve the integral. First, we'll square both sides, splitting the right hand side into two integrals with arbitrary variables, x and y .

$$J^2 = \left(\int_{x=-\infty}^{\infty} e^{-x^2} dx \right) \left(\int_{y=-\infty}^{\infty} e^{-y^2} dy \right)$$

Now, we need to combine them. We'll do that using Fubini's Theorem, turning the squared/multiplied integrals into a double integral.

$$J^2 = \int_{x=-\infty}^{\infty} \int_{y=-\infty}^{\infty} e^{-x^2} e^{-y^2} dy dx$$

Looking at this, the x and y covers the entire Cartesian Plane. So, we can compact the double integral to cover all real numbers on the 2D plane, \mathbb{R}^2 .

$$J^2 = \iint_{\mathbb{R}^2} e^{-x^2} e^{-y^2} dx dy = \iint_{\mathbb{R}^2} e^{-x^2-y^2} dx dy$$

The exponent on the e looks similar to the circle equation, so we're going to convert the integral into polar coordinates.

$$r^2 = x^2 + y^2 \quad dx dy = r dr d\theta$$

The angle only needs to go from $0 \rightarrow 2\pi$ to cover the entire graph, as well as $0 \rightarrow \infty$ for the radius.

$$J^2 = \int_{\theta=0}^{2\pi} \int_{r=0}^{\infty} e^{-r^2} r dr d\theta$$

Now, we can split this apart again by doing a reverse Fubini's Theorem and solving the left, simple, integral.

$$J^2 = \left(\int_{\theta=0}^{2\pi} d\theta \right) \left(\int_{r=0}^{\infty} e^{-r^2} r dr \right) = 2\pi \int_{r=0}^{\infty} e^{-r^2} r dr$$

From here, do u -sub letting $v = r^2$. We're using v because u is already used before in the problem.

$$\text{Let: } v = r^2, \quad dv = 2r dr$$

Substituting it in, the 2's and r 's cancel.

$$J^2 = \pi \int_{r=0}^{\infty} e^{-v} dv$$

Next, solve the integral and evaluate the bounds of integration.

$$J^2 = \pi \left[\frac{e^{-v}}{-1} \right]_{r=0}^{\infty} = \pi [0 - (-1)]$$

Finally, simplify in the brackets and take the square root of both sides, isolating J.

$$J^2 = \pi \rightarrow J = \sqrt{\pi}$$

Substituting back in the original function for J, we get:

$$\boxed{\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}}$$

Now, we've gone over the graph of the function as well as a relatively complex example. Let's do some problems!

$$\mathbf{1} \quad \frac{3}{2}!$$

$$\mathbf{2} \quad \int_0^{\infty} e^{-u} (x+1)^4 \, dx$$

$$\mathbf{3} \quad \int\limits_1^{\infty} \frac{(\ln(x))^3}{x^2} \, dx$$

$$4 \int\limits_0^{\infty} x^6 e^{-4x^2} \, dx$$

$$5 \int\limits_0^1 x^3 \sqrt{x \ln \left(\frac{1}{x}\right)} \, dx$$